

4. *Jako J, Fenyvesi A, Banai J*: Trombophilia in ulcerative colitis. *Orv Hetil* 2000; 141: 2139–2145.
5. *Socha P, Ryzko J, Janczyk W et al.*: Hepatic Vein Thrombosis as a complication of ulcerative colitis in a 12-year-old patient. *Dig Dis Sci* 2007; 52: 1293–1298.
6. *Lengle SJ, Nadler P, Jordan GW*: Arterial thrombosis in ulcerative colitis — transcatheter thrombolytic therapy. *West J Med* 1995; 162: 543–547.
7. *Hammer S, Meisner F, Dirschedl P et al.*: Procalcitonin: a new marker for diagnosis of acute rejection and bacterial infection in patients after heart and lung transplantation. *Transplant Immunology* 1998; 6: 235–241.
8. *Kareem E, Riem H, Firas S*: Inflammatory bowel disease-related thoracic Aortic Thrombosis. *South Med J* 2010; 103: 172–174. doi: 10.1097/SMJ.0b013e3181c95bc8.
9. *Miechlsler NP, Lator P, Valic E et al.*: Is inflammatory bowel disease an independent and disease specific risk factor for thromboembolism? *Gut* 2004; 53: 542–548.

**Corresponding author:**

*Małgorzata Malec-Milewska MD, PhD*  
*Department of Anaesthesiology and Intensive Care Medicine,*  
*Medical Centre for Postgraduate Education,*  
*Ul. Czerniakowska 231, 00–416 Warsaw, Poland*  
*e-mail: lmilewski@post.home.pl*

Anestezjologia Intensywna Terapia  
 2016, tom 48, numer 2, 153–156  
 ISSN 0209–1712  
 www.ait.viamedica.pl

## Ultrasound and fibreoptic-guided percutaneous tracheostomy in patient with deviated trachea

Kapil Dev Soni, Monica Jindal, Richa Aggarwal,  
 Mangilal Deganwa, Jagdish Prasad

*Department of Anaesthesia and Critical Care, All India Institute of Medical Sciences, New Delhi, India*

Sir,

A 78-year-old female, suffering from bronchial asthma with old pulmonary Koch's disease was admitted with a history of a fall followed by pain in the right hip. She was diagnosed with an inter-trochanteric fracture of the right femur. Her chest x-ray showed right middle zone and lower zone homogenous opacity with tracheal deviation (Fig. 1). She was admitted to the ICU after 4 days of hospital stay following acute breathlessness and arterial blood desaturation. The trachea was intubated and ventilatory support was commenced due to respiratory failure.

Due to predicted prolonged mechanical ventilation support, a percutaneous tracheostomy (PCT) was planned on the 5<sup>th</sup> day from ICU admission.

The trachea was located being shifted toward right side on palpation. A decision was made to use a USG for accurate localization of trachea and its real time puncture. Following skin disinfection, a vertical incision of 1 cm was performed 2 cm lateral to the midline at the level of the 2<sup>nd</sup> and 3<sup>rd</sup> tracheal cartilage (Fig. 2). The percutaneous tracheostomy (PCT) was performed with the Blue Rhino single dilator technique. The procedure was completed with minimal blood loss and no complications.



**Figure 1.** Chest x-ray of the patient showing tracheal deviation toward the right side



**Figure 2.** Modified incision for percutaneous tracheostomy 2 cm lateral to midline

Należy cytować wersję: *Soni KD, Jindal M, Aggarwal R, Deganwa M, Prasad J*: Ultrasound and fibreoptic-guided percutaneous tracheostomy in patient with deviated trachea. *Anaesthesiol Intensive Ther* 2016; 48: 148–149. doi 10.5603/AIT.2016.0025.

Percutaneous tracheostomy is the standard method of tracheostomy for patients in ICUs. However, the procedure is not without risks. The most common risks associated with PCT are as follows: haemorrhage, hypoxaemia, loss of airway, cannula misplacement, airway injury, pneumothorax, surgical emphysema, damage to the posterior tracheal wall and accidental decannulation in the immediate perioperative period [1].

Fibreoptic bronchoscopy can be used to identify the point of needle insertion into the trachea and for confirmation of the correct guide wire placement. This reduces the incidence of complications, especially posterior tracheal wall injury. Pre-procedure USG may be used to define the relevant airway anatomy, identify the tracheal midline, the blood vessels adjacent to the PCT insertion site, thereby decreasing the risk of bleeding. It also helps in estimating depth of the trachea from skin surface, as well as the tracheal diameter, thus assisting in tracheostomy tube size selection. Accurate placement of the needle into the trachea can be confirmed, while patients deemed difficult for PCT may also be identified using USG. However, its beneficial effects have not been established adequately in the literature. The use of USG increases the accuracy only marginally in centrally located trachea. A few studies have demonstrated an improvement in accuracy for PCT utilising USG. Rudas *et al.* [2] randomly assigned 50 patients to either traditional anatomical landmark or real time USG guidance. The latter method resulted in more accurate tracheal punctures.

However, the use of USG during the PCT in a patient with lung fibrosis, collapse and deviated trachea has rarely been described. Yavuz *et al.* [3], in their randomised controlled trial on 341 patients, reported the use of USG guidance for PCT in only one patient with deviated trachea due to lung fibrosis. To conclude, in difficult scenarios such as deviated trachea, the ability of USG to improve the accuracy of PCT increases considerably, along with the prevention of complications.

## ACKNOWLEDGEMENTS

1. The authors declare no financial disclosure.
2. The authors declare no conflict of interest.

## References:

1. Durbin CG: Early complications of tracheostomy. *Respir Care* 2005; 50: 511–15.
2. Rudas M, Seppelt I, Herkes R, Hislop R, Rajbhandari D, Weisbrodt L: Traditional landmark versus ultrasound guided tracheal puncture during percutaneous dilatational tracheostomy in adult intensive care patients: a randomised controlled trial. *Crit Care* 2014; 18: 514. doi: 10.1186/s13054-014-0514-0.
3. Yavuz A, Yilmaz M, Cemil G, Goya C, Alimoglu E, Kabaalioglu A: Advantages of US in percutaneous dilatational tracheostomy: randomized controlled trial and review of the literature. *Radiology* 2014; 273: 927–936. doi: 10.1148/radiol.14140088.

## Corresponding author:

Mangilal Deganwa, MD  
84B, Shanti Niwas, Arjun Nagar, New Delhi, India  
e-mail: mangilaldeganwa0606@gmail.com