

# Medical recommendations for home-confined footballers' training during the COVID-19 pandemic: from evidence to practical application

**AUTHORS:** Cristiano Eirale<sup>1</sup>, Giannicola Bisciotti<sup>1,2</sup>, Alessandro Corsini<sup>3</sup>, Christophe Baudot<sup>1</sup>, Gerard Saillant<sup>1</sup>, Hakim Chalabi<sup>1,2</sup>

<sup>1</sup> Paris Saint Germain FC, France

<sup>2</sup> Aspetar Sports and Orthopedics Hospital- Doha, Qatar

<sup>3</sup> Internazionale Milano FC, Italy

**ABSTRACT:** In early 2020, the world is facing a global emergency called COVID-19. Many professional footballers around the world are home confined. The maintenance of physical capacity is a fundamental requirement for the athlete, so the training sessions must be adapted to this unique situation. Specific recommendations must be followed concerning the type of training, its intensity, the precautions that have to be followed to avoid the possibility of contagion, and the restrictions in accordance with the presence of any symptoms. This article analyses the available scientific evidence in order to recommend a practical approach.

**CITATION:** Eirale C, Bisciotti G, Corsini A et al. Medical recommendations for home-confined footballers' training during the COVID-19 pandemic: from evidence to practical application. *Biol Sport*. 2020; 37(2):203–207.

Received: 2020-04-06; Reviewed: 2020-04-06; Re-submitted: 2020-04-10; Accepted: 2020-04-10; Published: 2020-04-10.

Corresponding author:

**Cristiano EIRALE**

Paris Saint Germain Football Club

53 avenue Emile Zola 92100

BOULOGNE-BILLANCOURT

E-mail: Ceirale@psg.fr

**Key words:**

Football

Soccer

Coronavirus

Training

Illness

## INTRODUCTION

In early 2020, the world is facing a global emergency called COVID-19 (coronavirus disease 2019). Many governments have requested the population to stay home unless necessary for proven reasons. Professional footballers have found themselves in a unique situation in which they were not only obliged to stop their professional activity but also to be confined in their houses.

### *The effect of detraining*

The maintenance of physical capacity is a fundamental requirement for the athlete. Specifically, for team sports athletes, maintaining a good level both of aerobic power and muscle strength is a fundamental prerequisite to preserve performance unchanged. The training physiological adaptation is a reversible process. Indeed, most aspects of physiological adaptation are lost during a prolonged period of inactivity [1]. The rate of loss is different for each physical capacity, being higher for endurance and strength endurance rather than speed and maximum strength. There is generally accepted to be an overall loss up to 10% of fitness for each week of total inactivity [2].

In professional football the annual season is subdivided into three periods: the pre-competition period (pre-season period), the competition period and the transition period (off-season) [3]. In professional football, the transition period rarely lasts more than one month [4]. In this period, there is an important reduction, or even a

complete stop, in football activity; therefore this forced pause of football activity during the pandemic of coronavirus can be considered a transition period. Obviously, it is very important that the players keep a good fitness level in order to be able to endure the rapid increase in training frequency and intensity, while trying, at the same time, to minimize the injury risk [5]. Indeed, many studies show that a loss of lean mass and muscle strength represents an important risk factor for muscle injuries in football [6–9]. Unfortunately, the off-season period generally has a negative influence on football players' body composition, increasing the percentage of fat mass and decreasing the percentage of lean mass or fat-free mass [4, 10–12]. Furthermore, as previously mentioned, a suspension or a substantial reduction of high-intensity aerobic activity for a period of 20 days or greater significantly decreases the  $VO_2$ max. For all these reasons, after the off-season period the players may show difficulty in tolerating high training volume and intensity in the first weeks of the pre-season period [4]. This is particularly evident in cases in which the changes in fat mass and free-fat mass are substantial [4]. With our current knowledge, it is impossible to predict what the real impact of the detraining period linked to the COVID-19 pandemic on the fitness status of athletes will be when they return to sport activity. There is no previous off-season period characterized by confinement. In any case, it is obvious that the level of fitness that players will

present when returning to normal training is closely linked to the following factors:

- i. Whether the player has contracted COVID-19 or not and, in case of contagion, if he/she shows any sequelae;
- ii. The duration of the detraining period and the confinement;
- iii. The level of physical activity that the player maintained during the detraining period.

Therefore, in accordance with the above points, it is extremely important that:

- at the time of resuming sports activity, a battery of tests allowing objective evaluation of the player's physical condition is performed;
- there is a suitable pre-competition period with a training programme including both aerobic training and strength conditioning activities [10,12].

### Training and the immune system

Indeed, physical activity may influence the response and the effectiveness of the immune system [13–18]. There is quite strong evidence that intense endurance sport activities, such as running, cycling, rowing or swimming, produce significant leukocytosis caused by an increase in numbers of B and T lymphocytes, neutrophils and NK cells in the systemic circulation [19,20]. Furthermore, acute severe exercise inducing an oxidative state results an acceleration of neutrophil apoptosis [21]. Therefore, at the end of demanding physical activity there is a drastic drop in circulating lymphocytes with consequent loss of efficiency of the immune system [17]. In exercise immunology a central dogma is that a strenuous exercise bout or a period of intense exercise impairs the effectiveness of the immune system leading to an “open window” of infection risk [22]. The concept that any kind of strenuous exercise can be considered ‘immunosuppressive’ has recently been challenged and the concept of an “open window” has been questioned [23,24]. However, the evidence provided for confuting the “open window” concept is not yet sufficiently convincing [25]. Thus, further relevant studies will be needed in the future. Therefore, considering both the high and dramatic specificity of the current pandemic period and the absence of evidence concerning sport activity during the COVID-19 pandemic [26], the training rules mentioned below are mainly based on the principle of “maximal caution” [27].

### Training intensity

Excessively intense training can weaken the immune system [22] and increase the risk of being contaminated by COVID-19 or of developing a serious form of it affecting the heart [28], the lung [29], the liver, the kidneys and the immune system [30]. For these reasons, we recommend avoiding intense training during the epidemic period.

### Training precautions

In countries where there are no limitations for outdoor sporting activities, outdoor running training is possible. The training must be

done individually and keeping the distance from other people imposed by the current health regulations. It is highly advisable to carry out the training in places not frequented by other people, preferring the times in which there is little flow of people. It is important to remember that the droplets containing COVID-19 travel a distance of 1.8 metres in the air and the average life of COVID-19 is 2.7 hours in the air, 13 hours on steel and 16 hours on polypropylene [31]. Furthermore, currently there is no scientific evidence regarding wearing protective masks in public spaces for asymptomatic persons [32]. For these reasons, it is highly inadvisable to perform any form of physical activity in public or private gyms frequented by several people. On the other hand, since there is a consensus in several countries to restrict social gathering, including gyms and meeting places, and to limit population movements [32], this recommendation may also be superfluous. Indeed, there are some sport activities such as yoga, Tai Ji Quan, and Qigong that can be preferred in this specific period and should be proposed to the population as a valid alternative to more popular sports [26]. Moreover, some alternative training methods such as software-generated partners should be proposed in order to maintain motivation [33].

### Cardio training

Suspension of high-intensity aerobic activity for 20 days or greater results in a significant decrease in  $VO_2\max$  [34,35]. This obviously can be a problem for a team sports athlete [36]; therefore, cardio training should be commenced as soon as possible. However, in compliance with the principle of maximal caution [27], it is highly recommended to observe the following rules:

- i. The duration of the cardio training sessions should not exceed 60 minutes [37]. It would be advisable to carry out two sessions of no more than 30 minutes during the day [38]. If the solution of two sessions is adopted, the latter must be interspersed with a recovery of at least three hours. It is very important to carry out suitable rehydration between the two training sessions [39].
- ii. The intensity of the effort should be limited to 80% of the maximum heart rate, corresponding to approximately 75-80% of the subject's maximum aerobic speed (i.e. the speed at which  $VO_2\max$  is reached) [40].
- iii. The cardio training can be performed both in the form of continuous running [41] and intermittent [42] or interval training [43], if the intensity of the effort indicated above is respected.

An average of 3-4 cardio training sessions performed by observing the following parameters are sufficient to maintain satisfactory functionality of both the aerobic and anaerobic lactic acid system [41], minimizing at the same time the possible risks associated with the COVID-19 emergency [26,28,44].

### Strength training

Strength training involves the use of equipment that usually is in public or private gyms. However, for the reasons of possible contagiousness explained above, attending public gyms is not recom-

mended. In the absence of suitable equipment, strength training can be carried out at home with elastic resistance [45] or body weight exercises, for example pull-ups, push-ups, sit-ups, dips, Nordic hamstring exercise, etc [46,47].

The strength training should respect the following rules:

- i. The duration of the strength training sessions should not exceed 60 minutes [37,48].
- ii. The use of maximum loads and exercises conducted at full muscle exhaustion (regardless of the load used) is not recommended [37,49]. Indeed, the lactate production typical of strength exercise conducted at complete muscle exhaustion promotes lymphocyte apoptosis [50].
- iii. During cross-fit sessions the intensity of the effort should be limited to 80% of the maximum heart rate [40].

Two weekly strength training sessions respecting the above-mentioned points are sufficient to maintain the strength characteristics in a well-trained athlete [51].

In any case, we do not recommend strength training to be carried out at maximal levels as there is evidence of a reduction of up to 15% of isometric strength in infected subjects who do not fully recover until a month after the illness [52]. Since in athletes the infection by coronavirus is often asymptomatic, athletes may potentially suffer from this kind of strength reduction even without a confirmed diagnosis of COVID infection, and therefore it is safer to train at submaximal levels.

### Stretching

Stretching exercise sufficient to maintain or to develop the range of motion should be included in the training sessions. The stretching exercises should involve the major muscle groups and be performed in a minimum of 2-3 sessions per week. Furthermore, stretching sessions should include both static and dynamic exercise [53].

### Training in case of fever or suspected infection

Fever is a normal physiological reaction of the body to an illness or immune stimulus assisting the immune system with mounting a response [54]. In case of fever, any type of physical activity must be suspended [55]. There is evidence from animals that strenuous exercise during an ongoing febrile infection can be dangerous, leading to an increased rate of complications and lethality [56]. Moreover, dehydration can contribute to hyperthermia [57]. In cases of febrile infections with systemic symptoms, recommendations on resuming physical activity vary from recommencement of sport activity once fever has resolved to waiting until 14 days after the symptom's resolution [56,58]. Moreover, the proximity with others during sport activity performed with an active pulmonary COVID 19 infection may potentially lead to spreading of the virus. Despite the lack of direct evidence of that, considering the method of transmission [44], it cannot be excluded.

In addition, in case of absence of fever, it is necessary to avoid

taking paracetamol as a preventive measure. Indeed, the effect of paracetamol could mask the onset of viral infection [59]. Some papers suggest a link between NSAIDs and both respiratory and cardiovascular adverse effects in several settings, but so far there is no evidence explicitly concerning COVID-19. Waiting for more robust evidence, we suggest a cautionary approach in their use [60]. Use of corticosteroids is also not advised because they increase the risk of infection, including viral infections [61].

For these reasons, we do not recommend any kind of training in case of fever following a COVID infection and, due to the lack of evidence, the principle of maximal prudence should be followed upon return to sport.

### Return to football

The day will come (we all hope soon) when COVID-19 will be only a memory. Looking forward to that day, as a final recommendation, we encourage the football medicine community both to apply maximal caution on the decision when to restart sport activity and to adopt a specific protocol to check for cardiological, pulmonary and, in general, systemic sequelae of COVID-19 in the athletes [27]. Since the geographical distribution of COVID-19 is unequal and the evolution of the pandemic is different in each country, the return to activity and the consequent programme of prevention should be tailored to each situation [62].

Moreover, the return to normal activity will have to be progressive and programmed. It is highly recommended that players return to sporting activity progressively, applying [63]:

- i) A preliminary phase normally performed during the confinement phase, during which the player restarts the training at home, with the training schedules sent and checked remotely by the club staff;
- ii. An individual training phase in which a few club staff members may assist the player during the pitch or gym training. During this phase, it is recommended that a maximum of two players can train on the pitch at the same time, keeping a safe distance between them;
- iii. A group training phase, with a maximum of 8 players at a time;
- iv. A collective training phase with full resumption of training for the whole group.

Furthermore, during the aforementioned phases, it will be necessary to respect a series of rigorous medical and hygienic-sanitary rules [63].

To date, there is no scientific evidence to demonstrate an ability of the pneumococcal vaccine (or any other vaccination) to protect against coronavirus infection.

Finally, at return to play there will probably be a congested period with 2-3 games every 7 days, after a period of detraining. Since both match congested periods [64] and activity load variation [65] have been associated with an increased risk of injury, attention should be paid to organizing for each player a tailored schedule of training and matches in order to avoid sudden variation of the load. This will also

be necessary in order to avoid potential exercise-induced depression of the immune system [66–68].

## CONCLUSION

During the 2020 COVID-19 pandemic, professional footballers are facing a unique situation. Maintaining physical fitness is important but the training should be modified accordingly in order to keep it safe and tailored to the new conditions. Therefore, it is important to maintain precise medical, hygienic-sanitary rules as well as to respect some physiological rules allowing, in this particular pandemic period,

safe training. Last, but certainly not least, the international medical community should rapidly establish a specific clinical protocol aimed at checking the athletes' medical status and physical fitness after the pandemic period caused by COVID-19. Therefore further studies are recommended in order to establish the evidence behind these protocols.

## Conflict of interest

All authors declare having NO conflict of interest

## REFERENCES

- Mallo J. Complex football. From Seirullos' structured training to frad's tactical periodization. Javier Mallo Sainz Editions; 2015.
- Varandas F, Medina D, Gomez A, Della Villa S. Late rehabilitation on the field. In: Injury and health problem in football. Springer Berlin Heidelberg; 2017. p. 571–9.
- Suarez-Arrones L, Lara-Lopez P, Maldonado R, Torreno N, De Hoyo M, Nakamura FY, et al. The effects of detraining and retraining periods on fat-mass and fat-free mass in elite male soccer players. *PeerJ*. 2019;
- Silva JR, Brito J, Akenhead R, Nassis GP. The Transition Period in Soccer: A Window of Opportunity. *Sports Med*. 2016;46(3):305–313.
- Gabbett TJ, Domrow N. Relationships between training load, injury, and fitness in sub-elite collision sport athletes. *J Sports Sci*. 2007;25(13):1507–1519.
- Croisier JL, Ganteaume S, Binet J, Genty M, Ferret JM. Strength imbalances and prevention of hamstring injury in professional soccer players: A prospective study. *Am J Sports Med*. 2008;36(8):1469–1475.
- De Hoyo M, Pozzo M, Sañudo B, Carrasco L, Gonzalo-Skok O, Domínguez-Cobo S, et al. Effects of a 10-week in-season eccentric-overload training program on muscle-injury prevention and performance in junior elite soccer players. *Int J Sports Physiol Perform*. 2015;10(1):46–52.
- Mendez-Villanueva A, Suarez-Arrones L, Rodas G, Fernandez-Gonzalo R, Tesch P, Linnehan R, et al. MRI-based regional muscle use during hamstring strengthening exercises in elite soccer players. *PLoS One*. 2016;11(9):e0161356.
- Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C, Opar DA. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): A prospective cohort study. *Br J Sports Med*. 2016;50(24):1524–1535.
- Koundourakis NE, Androulakis NE, Malliaraki N, Tsatsanis C, Venihaki M, Margioris AN. Discrepancy between exercise performance, body composition, and sex steroid response after a six-week detraining period in professional soccer players. *PLoS One*. 2014;9(2):e87803.
- Reinke S, Karhausen T, Doehner W, Taylor W, Hottenrott K, Duda GN, et al. The influence of recovery and training phases on body composition, peripheral vascular function and immune system of professional soccer players. *PLoS One*. 2009;4(3):1–7.
- Sotiropoulos A, Travlos AK, Gissis I, Souglis AG, Grezios A. The effect of a 4-week training regimen on body fat and aerobic capacity of professional soccer players during the transition period. *J Strength Cond Res*. 2009;23(6):1697–1703.
- Gleeson M. Immune system adaptation in elite athletes. *Curr Opin Clin Nutr Metab Care*. 2006;9(6):659–665.
- Brolinson PG, Elliott D. Exercise and the Immune System. *Clin Sports Med*. 2007;26(3):311–319.
- Nieman DC. Clinical implications of exercise immunology. *J Sport Health Sci*. 2012;(1)1:12–17.
- Lancaster GI, Febbraio MA. Exercise and the immune system: Implications for elite athletes and the general population. *Immunol Cell Biol*. 2016;94(2):115–6.
- Walsh NP, Oliver SJ. Exercise, immune function and respiratory infection: An update on the influence of training and environmental stress. *Immunol Cell Biol*. 2016;94(2):132–139.
- Nieman DC, Wentz LM. The compelling link between physical activity and the body's defense system. *J Sport Health Sci*. 2019;8(3):201–217.
- Pedersen BK, Hoffman-Goetz L. Exercise and the immune system: Regulation, integration, and adaptation. *Physiol Rev*. 2000;80(3):1055–1081.
- Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, et al. Position statement part one: Immune function and exercise. *Exerc Immunol Rev*. 2011;17:6–63.
- Syu G Da, Chen HI, Jen CJ. Severe exercise and exercise training exert opposite effects on human neutrophil apoptosis via altering the redox status. *PLoS One*. 2011;6(9):e24385.
- Simpson R, Campbell J, Gleeson M, Kruger K, Nieman DC, Pyne D, et al. Can exercise affect immune function to increase susceptibility to infection? *Exerc Immunol Rev*. 2020;26:8–22.
- Campbell JP, Turner JE. Debunking the myth of exercise-induced immune suppression: Redefining the impact of exercise on immunological health across the lifespan. *Front Immunol*. 2018;9:648.
- Campbell JP, Turner JE. There is limited existing evidence to support the common assumption that strenuous endurance exercise bouts impair immune competency. *Expert Rev Clin Immunol*. 2019;15(2):105–109.
- Zhu W. Should, and how can, exercise be done during a coronavirus outbreak? An interview with Dr. Jeffrey A. Woods. *J Sport Health Sci*. 2020;9(2):105–107.
- Chen P, Mao L, Nassis GP, Harmer P, Ainsworth BE, Li F. Wuhan coronavirus (2019-nCoV): The need to maintain regular physical activity while taking precautions. *J Sport Health Sci*. 2020;9(2):103–104.
- Corsini A, Bisciotti GN, Eirale C, Volpi P. Football cannot restart soon during the COVID-19 emergency! A critical perspective from the Italian experience and a call for action. *Br J Sports Med*. 2020;bjsports-2020-102306.
- Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020 Mar 5:1–2. doi: 10.1038/s41569-020-0360-5.
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507–13.
- Wang T, Du Z, Zhu F, Cao Z, An Y, Gao Y, et al. Comorbidities and multi-organ injuries in the treatment of COVID-19.

- The Lancet. 2020;395(10228):PE52
31. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect.* 2020;104(3):246–251.
  32. Flahault A. COVID-19 cacophony: is there any orchestra conductor? *Lancet.* 2020;395(10229):1037.
  33. Samendinger S, Hill CR, Kerr NL, Winn B, Ede A, Pivarnik JM, et al. Group dynamics motivation to increase exercise intensity with a virtual partner. *J Sport Health Sci.* 2019 May;8(3):289–297.
  34. Convertino VA. Cardiovascular consequences of bed rest: Effect on maximal oxygen uptake. *Med Sci Sports Exerc.* 1997;29(2):191–196.
  35. Bringard A, Pogliaghi S, Adami A, De Roia G, Lador F, Lucini D, et al. Cardiovascular determinants of maximal oxygen consumption in upright and supine posture at the end of prolonged bed rest in humans. *Respir Physiol Neurobiol.* 2010;172(1-2):53–62.
  36. Bisciotti GN, Volpi P, Alberti G, Aprato A, Artina M, Auci A, et al. Italian consensus statement (2020) on return to play after lower limb muscle injury in football (soccer). *BMJ Open Sport Exerc Med.* 2019;5(1):e000505.
  37. Hansen D, Niebauer J, Cornelissen V, Barna O, Neunhäuserer D, Stettler C, et al. Exercise Prescription in Patients with Different Combinations of Cardiovascular Disease Risk Factors: A Consensus Statement from the EXPERT Working Group. *Sports Med.* 2018 Aug;48(8):1781–1797.
  38. Hansen D, Dendale P, Coninx K, et al. The European Association of Preventive Cardiology Exercise Prescription in Everyday Practice and Rehabilitative Training (EXPERT) tool: A digital training and decision support system for optimized exercise prescription in cardiovascular disease. *Concept, definitions and construction methodology.* *Eur J Prev Cardiol.* 2017;24(10):1017–1031.
  39. Ranchordas MK, Dawson JT, Russell M. Practical nutritional recovery strategies for elite soccer players when limited time separates repeated matches. *J Int Soc Sports Nutr.* 2017;14:35.
  40. McMullen CW, Harrast MA, Baggish AL. Optimal running dose and cardiovascular risk. *Curr Sports Med Rep.* 2018;17(6):192–198.
  41. Dufour SP, Ponsot E, Zoll J, Doutreleau S, Lonsdorfer-Wolf E, Geny B, et al. Exercise training in normobaric hypoxia in endurance runners. I. Improvement in aerobic performance capacity. *J Appl Physiol (1985).* 2006 Apr;100(4):1238–48.
  42. Billat VL, Slawinski J, Bocquet V, Demarle A, Lafitte L, Chassaing P, et al. Intermittent runs at the velocity associated with maximal oxygen uptake enables subjects to remain at maximal oxygen uptake for a longer time than intense but submaximal runs. *Eur J Appl Physiol.* 2000;81(3):188–196.
  43. Laffite LP, Mille-Hamad L, Koralsztein JP, Billat VL. The Effects of Interval Training on Oxygen Pulse and Performance in Supra-threshold Runs. *Arch Physiol Biochem.* 2003;111(3):202–210
  44. Wang L-S, Wang Y-R, Ye D-W, Liu Q-Q. A review of the 2019 Novel Coronavirus (COVID-19) based on current evidence. *Int J Antimicrob Agents.* 2020 Mar 19:105948.
  45. Mascarin NC, De Lira CAB, Vancini RL, Pochini A de C, da Silva AC, Andrade M dos S. Strength training using elastic bands: Improvement of muscle power and throwing performance in young female handball players. *J Sport Rehabil.* 2017;26(3):245–252.
  46. Klika B, Jordan C. High-intensity circuit training using body weight: Maximum results with minimal investment. *ACSM's Health Fit J.* 2013;17(3):8–13.
  47. Seymore KD, Domire ZJ, DeVita P, Rider PM, Kulas AS. The effect of Nordic hamstring strength training on muscle architecture, stiffness, and strength. *Eur J Appl Physiol.* 2017;117(5):943–953.
  48. Loaza-Betancur AF, Bedoya EP, Dávila JM, Chulvi-Medrano I. Effect of Isometric Resistance Training on Blood Pressure Values in a Group of Normotensive Participants: A Systematic Review and Meta-analysis. *Sports Health.* 2020;1941738120908070.
  49. Westcott WL. Resistance training is medicine: Effects of strength training on health. *Curr Sports Med Rep.* 2012;11(4):209–16.
  50. Lagadic-Gossmann D, Huc L, Lecureur V. Alterations of intracellular pH homeostasis in apoptosis: Origins and roles. *Cell Death Differ.* 2004;11(9):953–961.
  51. Rønnestad BR, Hansen EA, Raastad T. In-season strength maintenance training increases well-trained cyclists' performance. *Eur J Appl Physiol.* 2010;110(6):1269–1282.
  52. Friman G. Effect of acute infectious disease on isometric muscle strength. *Scand J Clin Lab Invest.* 1977;37(4):303–8.
  53. Seirullo F. Sistemas dinámicos y rendimiento en deportes de equipo 1st meeting of complex system and sport. In Barcelona: INFEC; 2003.
  54. Broom M. Physiology of fever. *Paediatric nursing.* 2007.
  55. Russell J, Slutsky A. International consensus conferences in intensive care medicine: Ventilator-associated lung injury in ARDS. *Am J Respir Crit Care Med.* 1999;160(6):2118–24.
  56. Friman G, Wesslen L. Special feature for the Olympics: effects of exercise on the immunosystem: infections and exercise in high-performance athletes. *Immunol Cell Biol.* 2000 Oct;78(5):510–22.
  57. Trangmar SJ, González-Alonso J. Heat, Hydration and the Human Brain, Heart and Skeletal Muscles. *Sports Med.* 2019;49(Suppl 1):69–85.
  58. Rowsey PJ, Metzger BL, Carlson J, Gordon CJ. Long-term exercise training selectively alters serum cytokines involved in fever. *Biol Res Nurs.* 2009;10(4):374–380.
  59. Callaway E, Cyranoski D. China coronavirus: Six questions scientists are asking. *Nature.* 2020;577(7792):605–607.
  60. Little P. Non-steroidal anti-inflammatory drugs and covid-19. *BMJ.* 2020;368:m1185. Published 2020 Mar 27. doi:10.1136/bmj.m1185
  61. Dixon WG, Suissa S, Hudson M. The association between systemic glucocorticoid therapy and the risk of infection in patients with rheumatoid arthritis: Systematic review and meta-analyses. *Arthritis Res Ther.* 2011;13(4):R139.
  62. Eirale C, Gillogly S, Singh G, Chamari K. Injury & illness epidemiology in soccer - Effects of global geographical differences -A call for standardized & consistent research studies. *Biol Sport.* 2017;34(3):249–254.
  63. Liga de Fútbol Profesional. LaLiga Protocol regarding the resumption of training across LaLiga Clubs. 2020.
  64. Bengtsson H, Ekstrand J, Waldén M, Häggglund M. Muscle injury rate in professional football is higher in matches played within 5 days since the previous match: A 14-year prospective study with more than 130 000 match observations. *Br J Sports Med.* 2018;52(17):1116–1122.
  65. Malone S, Owen A, Newton M, Mendes B, Collins KD, Gabbett TJ. The acute:chronic workload ratio in relation to injury risk in professional soccer. *J Sci Med Sport.* 2017;20(6):561–565.
  66. Ferrari HG, Gobatto CA, Manchado-Gobatto FB. Training load, immune system, upper respiratory symptoms and performance in well-trained cyclists throughout a competitive season. *Biol Sport.* 2013;30(4):289–294.
  67. Trochimiak T, Hübner-Woźniak E. Effect of exercise on the level of immunoglobulin a in saliva. *Biol Sport.* 2012;29(4):255–261
  68. Moraes H, Aoki MS, Freitas CG, Arruda AFS, Drago G, Moreira A. IgA response and incidence of upper respiratory tract infections during intensified training in youth basketball players. *Biol Sport.* 2017;34(1):49–55.